

ON NUMERICAL COMPUTATIONAL SOLUTION OF SEVENTH ORDER BOUNDARY VALUE PROBLEMS

K. I. FALADE¹, I. G. BAKU, A. T. TIAMIYU AND I. ISYAKU

ABSTRACT: In this paper, we present and applied Exponentially Fitted Collocation Approximate Method for the numerical solution of seventh order boundary value problems. The approximate solution of the problem is computed using Maple 18 software after the problem was slightly perturbed and collocated. Three examples have been considered to illustrate the efficiency and implementation of the method and the results are compared with the exact solution and some existing work in literature.

Keywords and phrases: Exponentially Fitted Collocation Approximate Method, Seventh Order Boundary Value Problems, Maple 18 Software and Analytical solution

2019 Mathematical Subject Classification: A80

1. INTRODUCTION

The boundary value problems of higher order have been discussed due to their mathematical role as well as their potential for applications in hydromagnetic and hydrodynamic stability, modeling of induction motor behaviors and other fields of science and engineering. The application of seventh order BVPs available in engineering sciences. These problems arise in Mathematical modeling of induction motors with two rotor circuits [1]. Generally, this problem is difficult to solve analytically. Many authors have developed numerical techniques for the numerical solution of this form of boundary values problems. For instance, [2] contains theorems which detail the conditions for existence and uniqueness of solutions of higher boundary value problems. [3] applied Haar Wavelet approach for the solution of seventh order ordinary differential equations, moreover, [4] applied Differential transformation method to

Received by the editors October 21, 2018; Revised June 29, 2019 ; Accepted: July 30, 2020

www.nigerianmathematicalsociety.org; Journal available online at <https://ojs.ictp.it/jnms/>

¹Corresponding author

solve seventh-order boundary value problem and [5] proposed an efficient numerical solution for seventh order differential equation by using Septic B-Spline collocation method with non-uniform length proposed and just to mention a few. This paper is to employ an easy, fast and accurate numerical technique to obtain numerical solution of higher order boundary value problems and comparing results with other methods available in literature [6, 7, 8].

We consider general seventh order non-homogenous differential equation with mixed boundary conditions of the form:

$$\frac{d^7 y(t)}{dt^7} + a(t)y(t) = g(t) \quad ; \quad t \in [a, b] \quad (1)$$

subject to the boundary conditions;

$$\begin{aligned} y(t_0) &= \beta_1 \\ y'(t_0) &= \beta_2 \\ y''(t_0) &= \beta_3 \\ y'''(t_0) &= \beta_4 \\ y'(t_1) &= \beta_5 \\ y''(t_1) &= \beta_6 \\ y'''(t_1) &= \beta_7 \end{aligned} \quad (2)$$

where $a(t)$ and $g(t)$ are smooth functions and $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7$ are constants.

In this paper, we obtain numerical solution of equation (1) using exponentially fitted collocation approximate method. The whole idea of the method is to use power series as a basis function and its derivative substituted into a slightly perturbed equation which eventually collocate. However, use of Chebyshev Polynomials during the perturbation of the equations, which established in literature that Chebyshev Polynomials of the first kind possess properties that minimize errors

within the region of consideration [9]. The advantage of one of these properties called minimax

property is in this study harnessed in such a way that the derived formula is automated in symbolic algebraic program using MAPLE 18 software.

2. CONSTRUCTION OF COMPUTATIONAL TECHNIQUE

We formulate and employ exponentially fitted collocation approximate method to obtain numerical solution for seven order boundary value problem (1).

Suppose we have power series of the form;

$$y_N(t) = \sum_{k=0}^N s_k t^k \quad (3)$$

And Exponentially fitted approximate solution proposed [10].

$$y_N(t) \approx \sum_{k=0}^N s_k t^k + \tau_n e^t \quad (4)$$

where t represents the dependent variables in the problem, n represents the order of the dependent variable and $s_k y_N(t)$ ($k \geq 0$) are the unknown constants to be determined, N is the length of computation and degree of chebyshev polynomials defined [11].

Obtain seventh derivative of equation(3), we have;

$$y_N^7(t) = \sum_{k=7}^N k(k-1)(k-2)(k-3)(k-4)(k-5)(k-6)s_k t^{k-7} \quad (5)$$

substitute equations (3) and (5) into equation (1);

$$\left[\sum_{k=7}^N k(k-1)(k-2)(k-4)(k-5)(k-6)s_k t^{k-7} \right] + a(t) \left[\sum_{k=0}^N s_k t^k \right] = g(t)$$

$$\begin{aligned} & 5040s_7 + 40320ts_8 + 181440t^2s_9 + \dots \\ & + N(N-1)(N-2)(N-3)(N-4)(N-5)(N-6)t^{N-7}s_N \\ & + a(t)[s_0 + ts_1 + t^2s_2 + t^3s_3 + t^4s_4 + t^5s_5 + t^6s_6 + \dots + s_N t^N] \\ & = g(t) \end{aligned} \quad (6)$$

Collect the likes terms in equation (6), we have;

$$\begin{aligned} & a(t)s_0 + a(t)ts_1 + a(t)t^2s_2 + a(t)t^3s_3 + a(t)t^4s_4 + a(t)t^5s_5 + \\ & a(t)t^6s_6 + [5040 + a(t)t^7]s_7 + [40320t + a(t)t^8]s_8 + [181440t^2 + \\ & a(t)t^9]s_9 + \\ & \vdots \\ & + N(N-1)(N-2)(N-3)(N-4)(N-5)(N-6)t^{N-7}s_N \\ & + a(t)t^N = g(t) \end{aligned} \quad (7)$$

We perturbed and collocate equation 7, we have

$$\begin{aligned}
& a(t)s_0 + a(t)ts_1 + a(t)t^2s_2 + a(t)t^3s_3 + a(t_q)t_q^4s_4 + a(t_q)t_q^5s_5 + \\
& \quad a(t_q)t_q^6s_6 + [5040 + a(t_q)t_q^7]s_7 + [40320t_q + a(t_q)t_q^8]s_8 + [181440t_q^2 + \\
& \quad a(t_q)t_q^9]s_9 + \\
& \quad \vdots \\
& \quad + [N(N-1)(N-2)(N-3)(N-4)(N-5)(N-6)t_q^{N-7} \\
& \quad + a(t_q)t_q^N]s_N = g(t_q) + H(t_q)
\end{aligned} \tag{8}$$

where $t_q = a + \frac{(b-a)q}{N+2}$; $q = 1, 2, \dots, N+1$

$$\begin{aligned}
H(t_q) = & \tau_1 T_N(t_q) + \tau_2 T_{N-1}(t_q) + \tau_3 T_{N-2}(t_q) + \tau_4 T_{N-3}(t_q) + \tau_5 T_{N-4}(t_q) \\
& + \tau_6 T_{N-5}(t_q) + \tau_7 T_{N-6}(t_q)
\end{aligned}$$

Here $\tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \tau_6, \tau_7$ are free tau parameters to be determined and $T_N(t_q), T_{N-1}(t_q), T_{N-2}(t_q), T_{N-3}(t_q), T_{N-4}(t_q), T_{N-5}(t_q), T_{N-6}(t_q)$ are the Chebyshev Polynomials of degrees $N, N-1, N-2, N-3, \dots, N-6$ define in [11].

Equation (8) is called perturbed collocated seventh order equation. Coupled with boundary conditions (2) and using approximate solution (5), we have

$$\begin{aligned}
y(t_0) &= s_0 + \tau_7 e^0 = \beta_1 \\
y'(t_0) &= s_1 + \tau_7 e^0 = \beta_2 \\
y''(t_0) &= 2s_2 + \tau_7 e^0 = \beta_3 \\
y'''(t_0) &= 6s_3 + \tau_7 e^0 = \beta_4 \\
y'(t_1) &= s_0 + s_1 + s_2 + s_3 + s_4 + \dots + s_N + \tau_7 e^1 = \beta_5 \\
y''(t_1) &= s_1 + 2s_2 + 3s_3 + 4s_4 + \dots + Ns_N + \tau_7 e^1 = \beta_6 \\
y'''(t_1) &= 2s_2 + 6s_3 + 12s_4 + \dots + N(N-1)s_N + \tau_7 e^1 = \beta_7
\end{aligned} \tag{9}$$

where t_0 and $t_1 = 1$ Altogether, we obtained $(N+8)$ algebraic linear equations in $(N+8)$ unknown constants. Thus, we put the $(N+8)$ algebraic equations in Matrix form as:

$$YS = G \tag{10}$$

Here

$$Y = \begin{bmatrix} y_{11} & y_{12} & y_{13} & \dots & \dots & \dots & \dots & y_{1N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_1) \\ y_{21} & y_{22} & y_{23} & \dots & \dots & \dots & \dots & y_{2N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_2) \\ y_{31} & y_{32} & y_{33} & \dots & \dots & \dots & \dots & y_{3N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_3) \\ y_{41} & y_{42} & y_{43} & \dots & \dots & \dots & \dots & y_{2N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_4) \\ y_{51} & y_{52} & y_{53} & \dots & \dots & \dots & \dots & y_{2N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_5) \\ y_{61} & y_{62} & y_{63} & \dots & \dots & \dots & \dots & y_{2N} & T_N(t-1) & T_{N-1}(t_1) & \dots & \dots & \dots & \dots & T_{N-6}(t_6) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ y_{1N} & y_{2N} & y_{3N} & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 6 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \ell^1 \\ 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \dots & \dots & \dots & \dots & N & \ell^1 \\ 0 & 0 & 2 & 6 & 12 & 20 & 30 & 42 & 56 & \dots & \dots & \dots & \dots & N(N-1) & \ell^1 \end{bmatrix}$$

$$S = [s_0, s_1, s_2, \dots, s_N, \tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \tau_6, \tau_7]^T$$

$$G = [g(t_0), g(t_1), g(t_2), \dots, g(t_N), \beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7]^T$$

The unknown constants $s_0, s_1, s_2, s_3, s_4, \dots, s_N, \tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \tau_6$, and τ_7 are obtain using MAPLE 18 software which eventually substitute into the approximate solution (4).

3. NUMERICAL APPLICATION

In this section, we consider three numerical test problems to demonstrate the efficiency of the propose method. Their numerical solutions and absolute error were presented. The numerical results for each problem are presented in tabular forms and compared with the exact solutions and existing work in the literature [6, 7, 8].

Example 1

Consider non-homogeneous seventh order boundary value problem with constant coefficient[6].

$$\frac{d^7 y(t)}{dt^7} - ty(t) = e^t(t^2 - 2t - 6); \quad 0 \leq t \leq 1 \quad (11)$$

subject to boundary conditions;

$$\begin{aligned} y(0) &= 1, & y(1) &= 0 \\ y'(0) &= 0, & y'(1) &= -e \\ y''(0) &= -1, & y''(1) &= -2e \\ y'''(0) &= -2 \end{aligned} \quad (12)$$

The exact solution is

$$y(t) = (1 - t)e^t \quad (13)$$

EFCAM Technique

Compare equation (11) with equation (8), we have the following; $a(t) = t$, $g(t) = e^t(t^2 - 2t - 6)$ and taking computational length $N = 12$. We obtain;

$$\begin{aligned}
& -t_q s_0 - t_q^2 s_1 - t_q^3 s_2 - t_q^4 s_3 - t_q^5 s_4 - t_q^6 s_5 - t_q^7 s_6 + \\
& (-t_q^8 + 5040)s_7 + (-t_q^9 + 40320t_q)s_8 + (-t_q^{10} + 181440t_q^2)s_9 \\
& + (-t_q^{11} + 604800t_q^3)s_{10} + (-t_q^{12} + 1663200t_q^4)s_{11} + (-t_q^{13} \\
& + 3991680t_q^5)s_{12} - (8388608t_q^{12} - 50331648t_q^{11} + 132120576t_q^{10} \\
& - 199229440t_q^9 + 190513152t_q^8 - 120324096t_q^7 + 50692096t_q^6 \\
& - 14057472t_q^5 + 2471040t_q^4 - 25626t_q^3 + 13728t - q^2 - 288t_q + 1)\tau_1 \\
& - (2097152t - q^{11} - 11534336t - q^{10} + 27394048t_q^9 - 36765696t_q^8 \\
& + 30638080t_q^7 - 16400384t_q^6 + 5637632t_q^5 + 1208064t_q^4 + 151008t_q^3 \\
& - 9680t_q^2 + 242t_q - 1)\tau_2 - (52488t_q^{10} - 2621440t_q^9 + 5570560t_q^8 \\
& - 6553600t_q^7 + 4659200t_q^6 - 2050048t_q^5 + 549120t_q^4 - 84480t_q^3 \\
& + 6600t_q^2 - 200t_q + 1)\tau_3 - (131072t_q^9 - 589824t_q^8 + 1105920t_q^7 \\
& - 1118208t_q^6 + 658944t_q^5 - 228096t_q^4 + 44352t_q^3 - 4320t_q^2 \\
& + 162t_q - 1)\tau_4 - (32768t_q^8 - 131072t_q^7 + 212992t_q^6 - 180224t_q^5 \\
& - 84480t_q^4 - 21504t_q^3 + 2688t_q^2 - 128t_q + 1)\tau_5 \\
& - (8172t_q^7 - 28672t_q^6 + 39424t_q^5 - 26880t - q^4 \\
& + 9408t_q^3 - 1568t_q^2 + 98t_q - 1)\tau_6 \\
& - (2048t_q^6 - 6144t_q^5 + 6912t_q^4 \\
& - 3584t_q^3 + 640t_q^2 - 72t_q + 1)\tau_7 \\
& = e^{t_q}(t_q^2 - 2t_q - 6)
\end{aligned} \tag{14}$$

Collocate equation (14) as follows; $t_q = a + \frac{(b-a)q}{N+2}$; $q = 1, 2, \dots, N+1$, where $a = 0$, $b = 1$, $N = 12$

$$\begin{aligned}
t_1 &= \frac{1}{14}, t_2 = \frac{2}{14}, t_3 = \frac{3}{14}, t_4 = \frac{4}{14}, t_5 = \frac{5}{14}, \\
t_6 &= \frac{6}{14}, t_7 = \frac{7}{14}, t_8 = \frac{8}{14}, t_9 = \frac{9}{14}, t_{10} = \frac{10}{14}, \\
t_{11} &= \frac{11}{14}, t_{12} = \frac{12}{14}, t_{13} = \frac{13}{14},
\end{aligned}$$

Couple with boundary conditions (12) and equation (10), thus, we use MAPLE 18 software to obtain unknown constants of equations (14), we have

$$\begin{aligned}
S_0 &= 0.9999996125, S_1 = -0.00001551955341, \\
S_2 &= -0.5000077598,
\end{aligned}$$

$$\begin{aligned}
S_3 &= -0.33333359199, S_4 = -0.12500006494, \\
S_5 &= -0.0333333345403, \\
S_6 &= -0.006944474212, S_7 = -0.001190474683, \\
S_8 &= -0.000173633177, \\
S_9 &= -0.00002198698021, S_{10} = -0.00000255519516, \\
S_{11} &= -0.00000019988483, \\
S_{12} &= -0.00000004008267738, \tau_1 = 6.09759720510^{-8}, \\
\tau_2 &= 1.68735200510^{-9}, \\
\tau_3 &= 2.86328811910^{-7}, \tau_4 = -2.95003739310^{-9}, \\
\tau_5 &= 7.62366093610^{-7}, \tau_6 = 5.34546785410^{-7}, \\
\tau_7 &= 0.00001551955341
\end{aligned}$$

Substitute the above constants into approximate solution (4), we have;

$$\begin{aligned}
y_{12}(t) &\approx 0.9999996125 - 0.00001551955341t - 0.5000077598t^2 \\
&- 0.33333359199t^3 - 0.12500006494t^4 - 0.0333333345403t^5 \\
&- 0.006944474212t^6 - 0.001190474683t^7 - 0.000173633177t^8 \\
&- 0.00002198698021t^9 - 0.00000255519516t^{10} \\
&- 0.00000019988483t^{11} - 0.00000004008267738t^{12} \\
&+ 0.00001551955341e^t
\end{aligned} \tag{15}$$

Table 1: Numerical result for seventh order BVP Example 1

t	Exact Solution	Absolute Error EFCAM	Absolute Error [8]
0.0	1.0000000000	0.00	0.00
0.1	0.9946538262	1.10 E —19	9.00 E —10
0.2	0.9771222064	2.10 E —10	1.41 E —19
0.3	0.9449011656	4.10 E —10	4.87 E —09
0.4	0.8950948188	4.10 E —10	3.94 E —09
0.5	0.8243606355	3.10 E —10	9.28 E —10
0.6	0.7288475200	0.00	5.71 E —09
0.7	0.6041258121	0.00	6.24 E —08
0.8	0.4451081857	1.10 E —10	1.29 E —08
0.9	0.2459603111	0.00	1.05 E —08
1.0	0.0000000000	0.00	0.00

Example 2

Consider non-homogeneous seventh order boundary value problem with constant coefficient [7].

$$\frac{d^7 y(t)}{dt^7} + y(t) = -e^t(35 + 12t + t^2); \quad 0 \leq t \leq 1 \tag{16}$$

subject to boundary conditions;

$$\begin{aligned} y(0) &= 1, & y(1) &= 0 \\ y'(0) &= 1, & y'(1) &= -e \\ y''(0) &= 0, & y''(1) &= -4e \\ y'''(0) &= -3 \end{aligned} \tag{17}$$

The exact solution is

$$y(t) = t(1 - t)e^t \tag{18}$$

EFCAM Technique

Compare equation (16) with equation (8), we have the following; $a(t) = 1$, $g(t) = -e^t(35 + 12t + t^2)$ and taking computational length $N = 14$. We obtain;

$$\begin{aligned} & s_0 + t_q s_1 + t_q^2 s_2 + t_q^3 s_3 + t_q^4 s_4 + t_q^5 s_5 + t_q^6 s_6 + \\ & (t_q^7 + 5040)s_7 + (t_q^8 + 40320t_q)s_8 + (t_q^9 + 181440t_q^2)s_9 + \\ & + (t_q^{10} + 604800t_q^3)s_{10} + (t_q^{11} + 1663200t_q^4)s_{11} + (t_q^{12} + 3991680t_q^5)s_{12} \\ & + (t_q^{13} + 8648640t_q^6)s_{13} + (t_q^{14} + 17297280t_q^7)s_{14} \\ & - (134217728t_q^{14} - 939524096t_q^{13} + 2936012800t_q^{12} - 5402263552t_q^{11} + \\ & 6499598336t_q^{10} - 5369233408t_q^9 + 3111714816t_q^8 - 12710087680t_q^7 \\ & + 361181184t_q^6 - 69701632t_q^5 + 8712704t_q^4 \\ & - 65228t_q^3 + 25480t_q^2 - 392t_q + 1)\tau_1 \\ & - (33554432t_q^{13} - 218103808t_q^{12} + 627048448t_q^{11} - 1049624576t_q^{10} + \\ & 113311744t_q^9 - 825556992t_q^8 + 412778496t_q^7 - 141213696t_q^6 \\ & + 32361472t_q^5 - 4759040t_q^4 + 416416t_q^3 \\ & - (2097152t_q^{11} - 11534336t_q^{10} + 27394048t_q^9 - 36765696t_q^8 + \\ & 30638080t_q^7 - 16400384t_q^6 + 5637632t_q^5 - 1208064t_q^4 \\ & + 151008t_q^3 - 9680t_q^2 + 242t_q - 1)\tau_4 \\ & - (52488t_q^{10} - 2621440t_q^9 + 5570560t_q^8 - 6553600t_q^7 + 4659200t_q^6 \\ & - 2050048t_q^5 + 549120t_q^4 - 84480t_q^3 + 6600t_q^2 - 200t_q + 1)\tau_5 \\ & - (131072t_q^9 - 589824t_q^8 + 1105920t_q^7 - 1118208t_q^6 + \\ & 658944t_q^5 - 228096t_q^4 + 44352t_q^3 - 4320t_q^2 + 162t_q - 1)\tau_6 \\ & - (32768t_q^8 - 131072t_q^7 + 212992t_q^6 - 180224t_q^5 \\ & + 84480t_q^4 - 21504t_q^3 + 2688t_q^2 - 128t_q + 1)\tau_7 \\ & = e^{t_q}(35 + 12t_q + t_q^2) \end{aligned} \tag{19}$$

Collocate equation (19) as follows; $t_q = a + \frac{(b-a)q}{N+2}$; $q = 1, 2, \dots, N+1$, where $a = 0$, $b = 1$, $N = 14$

$$\begin{aligned} t_1 &= \frac{1}{16}, t_2 = \frac{2}{16}, t_3 = \frac{3}{16}, t_4 = \frac{4}{16}, t_5 = \frac{5}{16}, \\ t_6 &= \frac{6}{16}, t_7 = \frac{7}{16}, t_8 = \frac{8}{16}, t_9 = \frac{9}{16}, t_{10} = \frac{10}{16}, \\ t_{11} &= \frac{11}{16}, t_{12} = \frac{12}{16}, t_{13} = \frac{13}{16}, t_{14} = \frac{14}{16}, t_{15} = \frac{15}{16}, \end{aligned}$$

Couple with boundary conditions (17) and equation (10), thus, we use MAPLE 18 software to obtain unknown constants of equations (19), we have;

$$\begin{aligned} S_0 &= 0.00000228395394, S_1 = 1.000002284, \\ S_2 &= 0.000001141976973, \\ S_3 &= -0.4999996193, S_4 = -0.3333332368, \\ S_5 &= -0.1249999847, \\ S_6 &= -0.0333332822, S_7 = 0.006944433316, \\ S_8 &= -0.001190520128, \\ S_9 &= -0.0001735127129, S_{10} = -0.000022185107346, \\ S_{11} &= -0.0000023531718333, S_{12} = -0.000000323875814, \\ S_{13} &= 0.000000002083454712, S_{14} = -0.000000006118446529, \\ \tau_1 &= -0.00000008483126116, \tau_2 = 0.0000002398392547, \\ \tau_3 &= 0.0000004268330286, \tau_4 = 0.000001136805924, \\ \tau_5 &= -0.000001236660853, \tau_6 = 0.000002957827944, \\ \tau_7 &= -0.000002283953946 \end{aligned}$$

Substitute the above constants into approximate solution 4, we have;

$$\begin{aligned} y_{14}(t) &\approx 0.00000228395394 + 1.000002284t + 0.000001141976973t^2 \\ &- 0.4999996193t^3 - 0.3333332368t^4 - 0.1249999847t^5 \\ &- 0.0333332822t^6 + 0.006944433316t^7 - 0.001190520128t^8 \\ &- 0.0001735127129t^9 - 0.00002218510734t^{10} - 0.0000023531718t^{11} \\ &- 0.000000323875814t^{12} + 0.000000002083454712t^{13} \\ &- 0.000000006118446529t^{14} - 0.000002283953946e^t \end{aligned} \quad (20)$$

Table 2: Numerical result for seventh order BVP Example 2

t	Exact Solution	Absolute Error EFCAM	Absolute Error [7]
0.0	0.0000000000	0.00	0.00
0.1	0.0994653826	5.10 E —11	1.40 E —07
0.2	0.1954244413	1.10 E —10	6.40 E —07
0.3	0.2834703497	1.10 E —10	2.90 E —06
0.4	0.3580379275	0.00	4.40 E —06
0.5	0.41218031785	1.10 E —10	6.70 E —06
0.6	0.4373085120	2.10 E —10	6.40 E —06
0.7	0.4228880685	3.10 E —10	3.70 E —06
0.8	0.3560865485	2.10 E —10	3.30 E —07
0.9	0.2213642800	0.00	1.40 E —06
1.0	0.0000000000	2.10 E —10	

Not available

Example 3

Consider non-homogeneous seventh order boundary value problem with constant coefficient [8].

$$\frac{d^7 y(t)}{dt^7} - y(t) = -7e^t; \quad 0 \leq t \leq 1 \quad (21)$$

subject to boundary conditions;

$$\begin{aligned} y(0) &= 1, & y(1) &= 0 \\ y'(0) &= 0, & y'(1) &= -e \\ y''(0) &= -1, & y''(1) &= -2e \\ y'''(0) &= -2 \end{aligned} \quad (22)$$

The exact solution is

$$y(t) = (1 - t)e^t \quad (23)$$

EFCAM Technique

Compare equation (21) with equation (8), we have the following; $a(t) = -1$, $g(t) = -7e^t$ and taking computational length $N = 12$.

We obtain;

$$\begin{aligned}
& -s_0 - t_q s_1 - t_q^2 s_2 - t_q^3 s_3 - t_q^4 s_4 - t_q^5 s_5 - t_q^6 s_6 + \\
& (-t_q^7 + 5040)s_7 + (-t_q^8 + 40320t_q)s_8 + (-t_q^9 + 181440t_q^2)s_9 + \\
& + (-t_q^{10} + 604800t_q^3)s_{10} + (-t_q^{11} + 1663200t_q^4)s_{11} + (-t_q^{12}s_{12} \\
& + 3991680t_q^5) - (8388608t_q^{12} - 50331648t_q^{11} + 132120576t_q^{10} \\
& - 199229440t_q^9 + 190513152t_q^8 - 120324096t_q^7 + 50692096t_q^6 \\
& - 14057472t_q^5 + 2471040t_q^4 - 25626t_q^3 + 13728t - q^2 - 288t_q \\
& + 1)\tau_1 - (2097152t - q^{11} - 11534336t - q^{10} + 27394048t_q^9 \\
& - 36765696t_q^8 + 30638080t - q^7 - 16400384t_q^6 + 5637632t_q^5 \\
& + 1208064t_q^4 + 151008t_q^3 - 9680t_q^2 + 242t_q - 1)\tau_2 \\
& - (52488t_q^{10} - 2621440t_q^9 + 5570560t_q^8 - 6553600t_q^7 + \\
& 4659200t_q^6 - 2050048t_q^5 + 549120t_q^4 - 84480t_q^3 + \\
& 6600t_q^2 - 200t_q + 1)\tau_3 \\
& - (131072t_q^9 - 589824t_q^8 + 1105920t_q^7 - 1118208t_q^6 \\
& + 658944t_q^5 - 228096t_q^4 + 44352t_q^3 - 4320t_q^2 + 162t_q - 1)\tau_4 \\
& - (32768t_q^8 - 131072t_q^7 + 212992t_q^6 - 180224t_q^5 \\
& - 84480t_q^4 - 21504t_q^3 + 2688t_q^2 - 128t_q + 1)\tau_5 \\
& - (8172t_q^7 - 28672t_q^6 + 39424t_q^5 - 26880t - q^4 \\
& + 9408t_q^3 - 1568t_q^2 + 98t_q - 1)\tau_6 \\
& - (2048t_q^6 - 6144t_q^5 + 6912t_q^4 \\
& - 3584t_q^3 + 640t_q^2 - 72t_q + 1)\tau_7 \\
& = -7e^{t_q}
\end{aligned} \tag{24}$$

Collocate equation (24) as follows; $t_q = a + \frac{(b-a)q}{N+2}$; $q = 1, 2, \dots, N+1$, where $a = 0$, $b = 1$, $N = 12$

$$\begin{aligned}
t_1 &= \frac{1}{14}, t_2 = \frac{2}{14}, t_3 = \frac{3}{14}, t_4 = \frac{4}{14}, t_5 = \frac{5}{14}, \\
t_6 &= \frac{6}{14}, t_7 = \frac{7}{14}, t_8 = \frac{8}{14}, t_9 = \frac{9}{14}, t_{10} = \frac{10}{14}, \\
t_{11} &= \frac{11}{14}, t_{12} = \frac{12}{14}, t_{13} = \frac{13}{14}
\end{aligned}$$

Couple with boundary conditions (22) and equation (10), thus, we use MAPLE 18 software to obtain unknown constants of equations (23), we have;

$$\begin{aligned}
S_0 &= 0.9999823812, S_1 = -0.00001761881384, S_2 = -0.5000088094, \\
S_3 &= -0.3333362698, S_4 = -0.1250007342, S_5 = -0.03333347987, \\
S_6 &= -0.006944468155, S_7 = -0.001190479545,
\end{aligned}$$

$$\begin{aligned}
S_8 &= -0.0001736311935, S_9 = -0.00002198834869, \\
S_{10} &= -0.000002554945345, S_{11} = -0.0000001996845, \\
S_{12} &= -0.00000004015877, \tau_1 = 1.541463958 \times 10^{-7}, \\
\tau_2 &= 3.426494606 \times 10^{-8}, \tau_3 = 7.216480768 \times 10^{-7}, \\
\tau_4 &= 1.327787170 \times 10^{-7}, \tau_5 = 0.000001890892661, \\
\tau_6 &= 8.475452817 \times 10^{-7}, \tau_7 = 0.00001551955341
\end{aligned}$$

Substitute the above constants into approximate solution(4), we have;

$$\begin{aligned}
y_{12}(t) &\approx 0.9999823812 - 0.00001761881384t - 0.5000088094t^2 \\
&\quad - 0.3333362698t^3 - 0.1250007342t^4 - 0.03333347987t^5 \\
&\quad - 0.006944468155t^6 - 0.001190479545t^7 - 0.0001736311935t^8 \\
&\quad - 0.00002198834869t^9 - 0.000002554945345t^{10} \\
&\quad - 0.0000001996845t^{11} - 0.00000004015877t^{12} \\
&\quad + 0.00001551955341e^t.
\end{aligned} \tag{25}$$

Table 3: Numerical result for seventh order BVP Example 3

t	Exact Solution	Absolute Error EFCAM	Absolute Error [8]
0.0	1.0000000000	0.00	0.00
0.1	0.09946538262	1.10 E —10	1.10 E —16
0.2	0.9771222064	2.10 E —10	4.40 E —16
0.3	0.9449011656	2.10 E —10	2.20 E —14
0.4	0.8950948188	2.10 E —10	7.70 E —13
0.5	0.8243606355	0.00	1.10 E —11
0.6	0.7288475200	2.10 E —10	1.00 E —10
0.7	0.6041258121	1.10 E —10	6.40 E —10
0.8	0.4451081857	0.000	3.20 E —09
0.9	0.2459603111	1.10 E —10	1.30 E —09
1.0	0.0000000000	6.70 E —13	4.60 E —08




Figure 1. Exact Solution, EFCAM, [6]
Seventh order boundary value problem Example 1




Figure 2. Exact Solution, EFCAM, [7]
Seventh order boundary value problem Example 2




Figure 3. Exact Solution, EFCAM, [8]
Seventh order boundary value problem Example 3

4. CONCLUSION

In this paper, exponentially fitted collocation approximate technique has been introduced and applied to obtain numerical solutions of seventh order boundary value problems. To test the accuracy and efficiency of the proposed method, it has been tested on three examples of seventh order boundary value problems. It is found that the obtained results are giving little or no error comparing to exact solutions and existing numerical methods available in the literature. The advantage of this present method lies on utilization of MAPLE 18 software for all computational works which enhance the accuracy and efficiency.

REFERENCES

- [1] Richards G. and Sarma P.R.R., *Reduced order models for induction motors with two rotor circuits*, IEEE. *Trans. Ener. Conv.*, Vol. 9, pp. 673-678 DOI: **10.1109/60.368342**, 1994

- [2] R.P. Agarwal, *Boundary-value problems for higher order differential equations*, World Scientific, Singapore, 1986.
- [3] A. Padmanabha Reddy Manjula S. H, C. Sateesha and N. M. Bujurke, *Haar Wavelet Approach for the solution of seventh order ordinary differential equations*, IIETA Vol. 3, No. 2, pp. 108-114 DOI: **10.18280/mmep.030212**, 2016.
- [4] Shahid S. Siddiqi et-al, *Solution of Seventh Order Boundary Value Problem by Differential Transformation Method*, World Applied Sciences Journal 16 (11): 1521-1526, 2012 ISSN: **1818-4952**, 2012.
- [5] Y. Rajashekhar Reddy, *Efficient Numerical Solution for Seventh Order Differential Equation by using Septic B-Spline Collocation Method with non-uniform length Global*, Journal of Pure and Applied Mathematics.ISSN **0973-1768** Volume 13, Number 7 pp. 3543-3548, 2017
- [6] Kalyani, P. and Lemma, M.N., *Solutions of Seventh Order Boundary Value Problems Using Ninth Degree Spline Functions and Comparison with Eighth Degree Spline Solutions*, Journal of Applied Mathematics and Physics, **4**, 249-261, 2016.
- [7] Kasi viswanadham K.N. S and Reddy S.M, *Numerical solution of seventh order boundary value problems by petrov-galekin method with quintic B-splines as basis functions and septic B-splines as weight functions*, Int. Jl. Comp. Appl., vol. **122**, pp. 0975-8887, DOI: **10.5120/21700-481**, 2015.
- [8] Abdulla-Al-Mamun and Mohammad Asaduzzaman, *Solution of Seventh Order Boundary Value Problem by Using Variation Iteration Method* , International Journal of Mathematics and Computational Science Vol. **5**, No. 1,pp. 10-11, 2019.
- [9] Brever, K. S. and Eversion R. M., *On the errors incurred calculating derivatives using Chebyshev polynomials.*, Journal of Computer Physics, **99**, **56-67**, 1992.
- [10] Falade K.I (2015), *Exponentially fitted collocation approximation method for singular initial value problems and integro-differential equations.*, Ph.D Thesis (unpublished), University of Ilorin, Ilorin Nigeria 7-15, 2015.
- [11] Falade K.I, Tiamiyu A.T and Muhammad M.Y, *On Numerical Investigation and Assessment of variable Coefficient $C(t)$ on Airy differential equation*, Journal of National Association of Mathematical Physics Volume **48** pp36-37, 2018.

Department of Mathematics, Faculty of Computing and Mathematical Sciences, Kano University of Science and Technology, P.M.B 3244 Wudil, Kano State, Nigeria.
E-mail address: faladekazeem2016@kustwudil.edu.ng

Department of Mathematical Sciences, Federal University, Dutsin-ma P.M.B.5001, Dutsin-Ma, Katsina State, Nigeria.
E-mail address: ibaokun@fudutsinma.edu.ng

Department of Mathematics, Faculty of Computing and Mathematical Sciences, Kano University of Science and Technology, P.M.B 3244, Wudil, Kano State, Nigeria.
E-mail address: abdgafartunde@yahoo.com

Department of Mathematics, Faculty of Computing and Mathematical Sciences
 Kano University of Science and Technology, P.M.B 3244 Wudil Kano State Nigeria.
E-mail address: isyaku.idris@yahoo.com